

FREE BIASOCIATIVE GROUPOIDS¹

Snežana Ilić², Biljana Janeva³ Naum Celakoski⁴

Abstract. The subject of this paper is the study of the variety of groupoids that have the following property: each subgroupoid generated by two elements is a subsemigroup. A construction of free objects in this variety is given. Free objects in the variety of idempotent and commutative groupoids with the mentioned property are also constructed.

AMS Mathematics Subject Classification (1991): 03C05, 08B20

Key words and phrases: Biassociative groupoids, partial groupoids, partial semigroups, free groupoids, generating set, basis, base, exponent, primitive element.

0. Preliminaries

The idea of considering biassociative groupoids came out from [3], where monoassociative groupoids (i.e. groupoids with the property that each subgroupoid generated by one element is a subsemigroup) are investigated. The goal of this paper is a description of free objects in the varieties of groupoids with the property that each subgroupoid generated by a two-element set is a subsemigroup. In order to accomplish this, some definitions, notations and facts on free semigroups will be given below.

Let A be a nonempty set. Then the set of all finite (nonempty) sequences (a_1, a_2, \dots, a_n) , where $a_\nu \in A$, will be denoted by A^+ . The pair (A^+, \cdot) , where “ \cdot ” is the concatenation of sequences, is a free semigroup with the basis A . In the sequel, A^+ will denote the semigroup and its carrier, as well, and the element (a_1, a_2, \dots, a_n) of A^+ will be denoted simply by $a_1 a_2 \dots a_n$, or a^n in the case $a_1 = a_2 = \dots = a_n = a$.

The following propositions are true.

Proposition 0.1. *Let \mathcal{N} be the set of positive integers. Then:*

- (a) *The semigroup A^+ is cancellative.*
- (b) *For each $a \in A^+$ there is a unique pair $(b, k) \in A^+ \times \mathcal{N}$, such that $a = b^k$, where $b \neq c^r$, for any $c \in A^+$ and $r \in \mathcal{N} \setminus \{1\}$.*
- (c) *If $B \neq \emptyset$ and $B \subseteq C$, then $B^+ \subseteq C^+$.*
- (d) *$B \cap C \neq \emptyset \Rightarrow (B \cap C)^+ = B^+ \cap C^+$. \square*

¹This paper was partly supported by the Macedonian Academy of Sciences and Arts within the project "Free and Related to them Algebraic Structures". We are grateful to professor Ć. Ćupona, the manager of the project, for the ideas and permanent support.

²Faculty of Sciences and Mathematics, University of Niš, Niš, Serbia and Montenegro

³Faculty of Natural Sciences and Mathematics, University of Skopje, Skopje, Republic of Macedonia

⁴Faculty of Mechanical Engineering, University of Skopje, Skopje, Republic of Macedonia

In the assertion (b), b is called the *base* and k the *exponent* of a . An element $u \in A^+$ is said to be *primitive* in A^+ if and only if $(\forall v \in A^+, n \geq 2) (u \neq v^n)$. The notion of primitive element could be introduced for any semigroup S just substituting A^+ by S in the definition above.

A groupoid $\mathbf{G} = (G, \cdot)$ is said to be *biassociative* if and only if (shorter iff) for any $a, b \in G$, the subgroupoid S of \mathbf{G} generated by a and b , i.e. $S = \langle a, b \rangle$, is a subsemigroup of \mathbf{G} . Moreover, if S is commutative (idempotent, commutative and idempotent) subsemigroup of \mathbf{G} , then \mathbf{G} is said to be *commutative* (*idempotent*, *commutative idempotent*) *biassociative groupoid*, respectively. The class of all biassociative (commutative, idempotent, commutative and idempotent) groupoids will be denoted by *Bass* (*ComBass*, *IdBass*, *ComIdBass*), respectively.

Let $\mathbf{G} = (G, \cdot) \in \text{Bass}$ and $a, b \in G$. The subsemigroup C of \mathbf{G} , generated by a , i.e. $C = \langle a \rangle$, is described by $C = \{a^k \mid k \geq 1\}$. The subsemigroup S of \mathbf{G} generated by a, b , i.e. $S = \langle a, b \rangle$, in the case when $a \notin \langle b \rangle$ and $b \notin \langle a \rangle$ consists of all elements of the form $a^{\alpha_1} b^{\beta_1} \dots a^{\alpha_r} b^{\beta_r}$, where $\alpha_1, \beta_r \geq 0$, $\beta_1, \alpha_2, \dots, \beta_{r-1}, \alpha_r \geq 1$, and “ x^0 ” means “lack of any symbol”.

The class of biassociative groupoids is hereditary and closed under direct products and homomorphisms. Therefore:

Proposition 0.2. *The class of all biassociative groupoids is a variety.* \square

The following proposition is also true.

Proposition 0.3. *If $1 \leq |B| \leq 2$, then B^+ is a free object in *Bass* with the basis B .* \square

The corresponding proposition to 0.3 for *ComIdBass* is the following

Proposition 0.4. *If $|B| = 1$, then a free *ComIdBass* with the basis B is B itself. If $B = \{a, b\}$, $a \neq b$, then a free *ComIdBass* with the basis B is $\{a, b, ab\}$.* \square

Considering Proposition 0.3 (Proposition 0.4), we will give in Section 1 (Section 2) only the construction of a free groupoid in *Bass* (in *ComIdBass*) with a basis B , such that $|B| \geq 3$.

For this purpose we need some more definitions.

Let $G \neq \emptyset$, $D \subseteq G \times G$, and $\cdot : D \rightarrow G$ be a mapping. Then $\mathbf{G} = (G, D, \cdot)$ is called a *partial groupoid* with the *domain* D . A subset $P \subseteq G$ is said to be a *subgroupoid of the partial groupoid* \mathbf{G} iff

$$(a, b) \in P^2 \cap D \Rightarrow a \cdot b \in P.$$

A subgroupoid of a partial groupoid need not be a groupoid, but it is a partial groupoid with the domain $P^2 \cap D$.

Let $\mathbf{S} = (S, D, \cdot)$ be a partial groupoid. \mathbf{S} is called a *partial semigroup*⁵ iff

$$(1) \quad (\forall a, b, c \in S)((ab)c, a(bc) \in S \Rightarrow (ab)c = a(bc)).$$

Let P be a subgroupoid of a partial groupoid \mathbf{G} . If \mathbf{P} is a partial semigroup, then \mathbf{P} is called a *partial subsemigroup* of \mathbf{G} .

A partial groupoid $\mathbf{G} = (G, D, \cdot)$ is said to be a *partial commutative (idempotent, commutative idempotent) groupoid* iff

$$(\forall a, b \in G)(ab \in G \Rightarrow ba \in G \wedge ab = ba),$$

$$((\forall a \in G)(a^2 \in G \Rightarrow a = a^2),$$

$$(\forall a, b \in G)(ab, a^2 \in G \Rightarrow ba \in G \wedge ab = ba \wedge a^2 = a)),$$

respectively.

The following proposition is also true.

Proposition 0.5. *Let K, P be subgroupoids of the partial groupoid $\mathbf{G} = (G, D, \cdot)$. If $K \cap P \neq \emptyset$, then $K \cap P$ is a subgroupoid of \mathbf{G} . \square*

Let \mathbf{G} be a partial groupoid, $\emptyset \neq A \subseteq G$, $\{P_i \mid i \in I\}$ the family of all subgroupoids of \mathbf{G} containing A , and $P = \bigcap_{i \in I} P_i$. Then $P \neq \emptyset$, and (by Proposition 0.5) P is a subgroupoid of \mathbf{G} which is called the *subgroupoid of \mathbf{G} generated by A* and is denoted by $P = \langle A \rangle$.

If \mathbf{G} and \mathbf{G}' are partial groupoids and $\varphi : G \rightarrow G'$ is a mapping, then φ is called a *partial homomorphism* from \mathbf{G} into \mathbf{G}' iff

$$(2) \quad (\forall x, y \in G)(xy \in G, \varphi(x)\varphi(y) \in G' \Rightarrow \varphi(xy) = \varphi(x)\varphi(y)).$$

Using the notions of subgroupoid of a partial groupoid generated by a non-empty set and partial homomorphism, one can define a partial free object in a class of partial groupoids in a usual way.

In order to give constructions of free objects in the varieties *Bass* and *ComIdBass* we need definitions of a partial biassociative groupoid and a free partial biassociative groupoid.

A partial groupoid $\mathbf{G} = (G, D, \cdot)$ is said to be *partial biassociative groupoid* (or partial *Bass*-groupoid) iff for any $a, b \in G$, $\langle a, b \rangle$ is a partial subsemigroup of \mathbf{G} .

A partial *Bass*-groupoid \mathbf{H} is said to be a *free partial Bass-groupoid with the basis B* ($B \neq \emptyset$), if \mathbf{H} is generated by B and if $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$ is a mapping, then there is a (unique) mapping $\varphi : H \rightarrow G$, such that φ is a partial homomorphism that is an extension of λ .

⁵A partial semigroup $\mathbf{S} = (S, D, \cdot)$ could be defined as follows

$$(\forall a, b, c \in S)((ab)c \in S \Rightarrow a(bc) \in S \wedge (ab)c = a(bc)),$$

but in this paper we will consider the one satisfying (1).

1. Construction of a free biassociative groupoid

The construction of a free biassociative groupoid with a given basis B will be given only for $|B| \geq 3$, as it was mentioned in Section 0. It will be given in several steps. In fact, an inductive construction of a chain $H_0, H_1, \dots, H_k, \dots$ of partial biassociative groupoids will be given such that its union will be a free object in *Bass* with the basis B .

The first step will be the construction of H_1 . To make the reading easier, we give the full construction when $|B| = 3$, $B = \{a, b, c\}$, and then we give just a short note for the case $|B| > 3$. Some auxiliary assertions in this section will be marked as 1.x.x.

1.1. Construction of H_1

The set $B = \{a, b, c\}$ has no structure, so it is assumed that $H_0 = B$ is a partial groupoid with the domain $D_0 = \emptyset$. Define the set H_1 by:

$$H_1 = \{a, b\}^+ \cup \{a, c\}^+ \cup \{b, c\}^+$$

(or, in general, $H_1 = \bigcup \{\{x, y\}^+ \mid x, y \in H_0, x \neq y\}$).

The fact that H_1 is a union of infinite sets, each being a free semigroup with a two-element basis, implies that:

1.1.1. $\mathbf{H}_1 = (H_1, D_1, \cdot)$ is a partial groupoid with the domain

$$D_1 = \{(t, u) \mid \{t, u\} \subseteq \{a, b\}^+ \vee \{t, u\} \subseteq \{a, c\}^+ \vee \{t, u\} \subseteq \{b, c\}^+\},$$

(or, in general, $D_1 = \bigcup \{(\{x, y\}^+)^2 \mid x, y \in H_0, x \neq y\}$). \square

Note that H_1 is a union (in general not disjoint) of free semigroups. It is not a groupoid, in the case $|B| \geq 3$. For example, if $a, b, c \in B$, $a \neq b \neq c \neq a$, then $ab, bc \in H_1$, but $(ab, bc) \notin D_1$, i.e. the "product" $ab \cdot bc$ does not exist in \mathbf{H}_1 . The elements of B are primitive elements in \mathbf{H}_1 , but there are others, such as ab, bc, \dots .

We give below some properties of \mathbf{H}_1 .

1.1.2. \mathbf{H}_1 is a partial Bass-groupoid and

$$x, y \in H_1 \Rightarrow ((x, y) \in D_1 \iff (y, x) \in D_1). \square$$

The next proposition is true for H_1 , but not for $H_k, k \geq 2$.

1.1.3. If $x, y, z \in H_1$, then $x(yz) \in H_1 \Rightarrow (xy)z \in H_1$, and in this case, $x(yz) = (xy)z$. \square

1.1.4. \mathbf{H}_1 is a free partial Bass-groupoid with the basis B .

Proof. Clearly, B generates \mathbf{H}_1 . Let $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$ be a mapping. If $(x, y) \in D_1$, then $x, y \in \{u, v\}^+$, where $u, v \in B = \{a, b, c\}$. Since $\{u, v\}^+$ is a free semigroup with the basis $\{u, v\}$, then there is a homomorphic extension ψ_1 of λ_1 from $\{u, v\}^+$ into \mathbf{G} , where λ_1 is the restriction of λ on the set $\{u, v\}$. We put $\varphi_1(xy) = \psi_1(xy) = \psi_1(x)\psi_1(y) = \varphi_1(x)\varphi_1(y)$. It is clear that φ_1 is a partial homomorphism from \mathbf{H}_1 into \mathbf{G} . \square

1.2. Construction of \mathbf{H}_2

Many “products” of elements of H_1 are not defined in H_1 , such as $a \cdot (bc)$, $b \cdot (ac)$, $(ab) \cdot (ac)$. To provide their existence, we extend H_1 to H_2 as follows:

$$H_2 = H_1 \cup (\cup\{\{t, u\}^+ \mid t, u \text{ are primitive elements in } H_1 \ \& \ (t, u) \notin D_1\}).$$

Remark 1. In definition to H_2 we could have taken the union of the collection $\{\{v, w\}^+ \mid v, w \in H_1, (v, w) \notin D_1\}$, for if v, w are not primitive elements in H_1 , then $v = t^m$, $w = u^n$ for some $t, u \in H_1$, and $\{v, w\}^+ \subseteq \{t, u\}^+$.

Remark 2. Denote $C_1 = \cup\{\{t, u\}^+ \mid t, u \text{ are primitive elements in } H_1 \ \& \ (t, u) \notin D_1\}$. Then: $H_1 \cap C_1 = \{v^n \mid v \text{ is a primitive element in } H_1, n \geq 1\} \neq \emptyset$, $C_1 \setminus H_1$ is infinite. For example, the set $\cup\{t \cdot u \mid t, u \text{ are primitive elements in } H_1 \ \& \ (t, u) \notin D_1\}$ is a proper subset of $C_1 \setminus H_1$.

Remark 3. If $v, w \in H_1$, then $v \cdot w$ is defined in H_2 iff $v \cdot w$ is defined in H_1 or $v \cdot w \in \{t, u\}^+$ for some primitive elements $t, u \in H_1$, such that $(t, u) \notin D_1$.

Remark 4. If t, u, v are primitive elements in H_1 such that $tu, uv \notin H_1$, then $(tu) \cdot v \notin H_2$ or $t \cdot (uv) \notin H_2$.

1.2.1. \mathbf{H}_2 is a partial groupoid with the domain

$$(3)D_2 = D_1 \cup (\cup\{\{t, u\}^+ \mid t, u \text{ are primitive elements in } H_1 \ \& \ (t, u) \notin D_1\}).$$

and $H_1^2 \subset D_2$.⁶ \square

Note that the union in (3) need not be disjoint. Some properties of H_2 will be listed bellow.

1.2.2. Each element in H_2 has a uniquely determined base and exponent. \square

1.2.3. \mathbf{H}_2 is a partial biassociative groupoid. \square

Note that \mathbf{H}_2 is not a partial semigroup, as $(ab)c \neq a(bc)$, although $(ab)c, a(bc) \in H_2$.

1.2.4. If $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$ is a mapping, then there is a unique partial homomorphism $\varphi_2 : \mathbf{H}_2 \rightarrow \mathbf{G}$, such that φ_1 is the restriction of φ_2 on the set H_1 .

Proof. Let $\mathbf{G} \in \text{Bass}$, and $\lambda : B \rightarrow G$ be a mapping. Then $\varphi_1 : H_1 \rightarrow G$ is a partial homomorphism defined as in the proof of 1.1.4. If $x, y \in H_2$, $(x, y) \in D_2$ and $x, y \in \{u, v\}^+$, where u, v are primitive elements in H_1 , then φ_2 is defined in the same way as φ_1 in 1.1.4. \square

⁶ $A \subset B$ iff $A \subseteq B$ and $A \neq B$.

1.3. Construction of H_n ($n \geq 3$)

Assume that the partial Bass groupoids $B = H_0, H_1, \dots, H_k$ are defined and the following conditions are satisfied:

a) For each i , $0 \leq i \leq k$, $H_i^2 \subset D_{i+1}$.

b) For each $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$, there is a chain of partial homomorphisms $\lambda = \varphi_0 \subseteq \varphi_1 \subseteq \dots \subseteq \varphi_{k+1} \subseteq \dots$, where $\varphi_k : H_k \rightarrow G$ for any $k \geq 0$.

Now, define H_{k+1} in the same way as H_2 :

$$H_{k+1} = H_k \cup (\cup\{\{t, u\}^+ \mid t, u \text{ are primitive elements in } H_k \ \& \ (t, u) \notin D_k\}).$$

1.3.1. H_{k+1} is a partial Bass-groupoid with the domain

$$D_{k+1} = D_k \cup (\cup\{\{t, u\}^+{}^2 \mid t, u \text{ are primitive elements in } H_k \ \& \ (t, u) \notin D_k\}).$$

□

Note that

$$D_{k+1} = H_k^2 \cup (\cup\{\{t, u\}^+{}^2 \mid t, u \text{ are primitive elements in } H_k \ \& \ (t, u) \notin D_k\}).$$

1.3.2. $(\forall k \geq 0) (H_k^2 \subset D_{k+1} \text{ and } D_k \subset H_k^2)$.

Proof. The proof will be given by induction on k for both statements at the same time.

Recall that $H_0 = B$, $D_0 = \emptyset$ and $D_1 = \cup\{\{x, y\}^+{}^2 \mid x, y \in H_0, x \neq y\}$. Clearly, $D_0 \subset H_0^2$, and $((ab), b) \in D_1$, but $((ab), b) \notin H_0^2$, i.e. $H_0^2 \subset D_1$. Thus 1.3.2 is true for $k = 0$.

We also give the proof for $k = 1$, i.e. $H_1^2 \subset D_2$ and $D_1 \subset H_1^2$.

Since $H_1 = \{a, b\}^+ \cup \{a, c\}^+ \cup \{b, c\}^+$, it follows that $(ab, c) \in H_1^2$, but $(ab, c) \notin D_1$, and thus $D_1 \subset H_1^2$. It is easily seen that there are elements $x, y, u \in H_1 \setminus H_0$, such that $(x, y) \notin D_1$, and $u \in \{x, y\}^+$ (for example: $x = ab$, $y = ac$, $u = (ab)^2$ are in $H_1 \setminus H_0$, $(ab, ac) \notin D_1$ and $(ab)^2 \in \{ab, ac\}^+$). Then $(xy, u) \notin H_1^2$, but $(xy, u) \in D_2$, i.e. 1.3.2 is true for $k = 1$.

Suppose that $H_r^2 \subset D_{r+1}$, and $D_r \subset H_r^2$, for each $r \in \{0, 1, \dots, k\}$, $k > 0$. We will prove that

$$H_{k+1}^2 \subset D_{k+2} \text{ and } D_{k+1} \subset H_{k+1}^2.$$

By the inductive hypothesis and the definitions of H_r, D_r , we have that $H_k \subset H_{k+1}$ and there are $x, y, u \in H_{k+1} \setminus H_k$, such that $(x, y) \notin D_k$ (as $D_k \subset H_k^2$) and $u \in \{x, y\}^+$. Then $(xy, u) \notin H_{k+1}^2$, but $(xy, u) \in D_{k+2}$. If $x, y, u \in H_{k+1} \setminus H_k$ are different primitive elements such that $u \notin \{x, y\}^+$, then $xy, u \in H_{k+1}$, $(xy, u) \in H_{k+1}^2$, but $(xy, u) \notin D_{k+1}$. Thus, $D_{k+1} \subset H_{k+1}^2$. □

1.3.3. Each element in H_{k+1} has a unique base and exponent. □

1.3.4. Let $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$ be a mapping. Then there is a unique partial homomorphism $\varphi_{k+1} : H_{k+1} \rightarrow G$, such that φ_k is the restriction of φ_{k+1} on H_k .

Proof. Let $(x, y) \in D_{k+1}$. If $(x, y) \in D_{k+1} \cap H_k^2$, then $\varphi_{k+1}(xy) = \varphi_k(x)\varphi_k(y)$. If $(x, y) \in D_{k+1} \setminus H_k^2$, then $x, y \in \{u, v\}^+$, for some primitive elements $u, v \in H_k$, such that $(u, v) \notin D_k$. Thus, $xy = u^{\alpha_1}v^{\beta_1} \dots u^{\alpha_r}v^{\beta_r}$, and we define

$$\varphi_{k+1}(xy) = \varphi_k(u)^{\alpha_1} \varphi_k(v)^{\beta_1} \dots \varphi_k(v)^{\beta_r}.$$

It is clear that φ_{k+1} is a partial homomorphism, and φ_k is the restriction of φ_{k+1} on H_k . \square

Theorem 1. *If $H = \bigcup_{k \geq 0} H_k$, then \mathbf{H} is a free biassociative groupoid with the basis B .*

Proof. First, let $x, y \in H$. Then there is a $k \in \mathcal{N}$, such that $x, y \in H_k$ and by 1.3.2, $(x, y) \in D_{k+1}$. Thus $x \cdot y \in H_{k+1} \subseteq H$, i.e. \mathbf{H} is a groupoid. Now, we will prove that $\mathbf{H} \in \text{Bass}$. Let $x, y \in H$, i.e. there is a k , such that $(x, y) \in D_k$. Then $\langle x, y \rangle$ is a subgroupoid of \mathbf{H} . Let $u, v, w \in \langle x, y \rangle$. Then $(u, v), (uv, w), (v, w), (u, vw) \in D_s$, for some $s \geq k$. As H_k is a partial Bass-groupoid for each k , it follows that $(uv)w = u(vw) \in H_s \subseteq H$. Thus, $\langle x, y \rangle$ is a subsemigroup, i.e. $\mathbf{H} \in \text{Bass}$. Let $\mathbf{G} \in \text{Bass}$ and $\lambda : B \rightarrow G$ be a mapping. Define $\varphi : H \rightarrow G$ as follows. If $(x, y) \in D_k$, then $\varphi(xy) = \varphi_k(x)\varphi_k(y)$. It is clear that φ is a homomorphism, such that $\varphi_0 = \lambda$ is the restriction of φ on the set B . (Note that, by the construction, B generates \mathbf{H} .) \square

Remark 5. If we consider the class of *ComBass*, then Theorem 1 can be restated for *ComBass* by adding commutativity. The construction of free commutative biassociative groupoid with a given basis B is essentially the same, except that it is based on a free commutative semigroup generated by two elements a and b , i.e. $\{a, b\}^{(+)}$ instead on a free semigroup $\{a, b\}^+$.

Moreover, the following statements for \mathbf{H}_k are also true, for each $k \in \mathcal{N}$.

1.3.5. *If $x, y \in H_k$, then $(x, y) \in D_k$ iff $(y, x) \in D_k$, and $\langle x, y \rangle$ is a subsemigroup of H_k .* \square

1.3.6 *qbt \mathbf{H}_k is a cancellative partial groupoid, i.e.*

$$(x, y), (x, z) \in D_k \Rightarrow (xy = xz \Rightarrow y = z), \text{ and}$$

$$(x, z), (y, z) \in D_k \Rightarrow (xz = yz \Rightarrow x = y).$$

Proof. \mathbf{H}_1 is a cancellative groupoid. Let the statement be true for all \mathbf{H}_r , $r \leq k$, and let $(x, y), (x, z) \in D_{k+1} \setminus H_k^2$ and $xy = xz$. Then $x, y \in \{u, v\}^+$, for some primitive elements $u, v \in H_k$ such that $(u, v) \notin D_k$ and $xy = xz \in \{u, v\}^+$. As $\{u, v\}^+$ is a free semigroup generated by $\{u, v\}$, it is a cancellative semigroup, and thus $y = z$. \square

2. Construction of Free Commutative Idempotent Biassociative Groupoids

We will consider here the class of commutative idempotent biassociative groupoids (*ComIdBass*) defined in Section 0. Clearly, if $\mathbf{G} \in \text{ComIdBass}$, then $\mathbf{G} \in \text{Bass}$ and \mathbf{G} is commutative and idempotent groupoid. Considering Proposition 0.5, we obtain that:

$$\mathbf{G} \in \text{ComIdBass} \iff (\forall x, y \in G)(x, y) = \{x, y, xy\},$$

where $xy = yx$.

Let us note that the following is valid:

Proposition 2.1 *If a, b are different objects, then the groupoid $\mathbf{H} = (\{a, b, ab\}; \cdot)$ defined by*

\cdot	a	b	ab
a	a	ab	ab
b	ab	b	ab
ab	ab	ab	ab

is a free semilattice with the basis $\{a, b\}$. \square

We will consider the case $|B| = 3$. The case $|B| > 3$ will not be considered, as the construction of a free *ComIdBass*-groupoid with the basis B , is essentially the same as in the case $|B| = 3$.

Let $B = \{a, b, c\}$, $a \neq b \neq c \neq a$. We will construct a chain $H_0, H_1, \dots, H_k, \dots$ of partial *ComIdBass*-groupoids by induction on k .

Define $H_0 = B$ and a partial order \leq_0 by: $a <_0 b <_0 c$. H_0 is a partial *ComIdBass* groupoid with the domain $D_0 = \emptyset$. Put $H_1 = H_0 \cup \{ab, ac, bc\}$, and define \leq_1 to be the lexicographic order on H_1 generated by \leq_0 . Then $\mathbf{H}_1 = (H_1, \cdot)$ is a partial *ComIdBass* groupoid with the domain

$$D_1 = \{(x, y) \mid x, y \in H_0\} = H_0^2.$$

Suppose that \mathbf{H}_k and \leq_k are defined such that \mathbf{H}_k is a partial *ComIdBass*-groupoid. Define

$$(4) \quad H_{k+1} = H_k \cup \{x(yz) \mid x, yz \in H_k, x <_k yz, x \neq y, x \neq z, x \neq yz\}$$

and \leq_{k+1} to be the lexicographic order on H_{k+1} generated by \leq_k .

Proposition 2.2. *\mathbf{H}_k is a partial *ComIdBass*-groupoid, for any $k \in \mathcal{N}$, with the domain $D_k = \{(x, y) \mid x, y \in H_{k-1}\} = H_{k-1}^2$.*

Proof. \mathbf{H}_0 and \mathbf{H}_1 are partial *ComIdBass* groupoids. Assume that \mathbf{H}_k is a partial *ComIdBass* groupoid, and consider H_{k+1} defined by (4).

If $u, v \in H_{k+1}$, $(u, v) \in D_{k+1}$, then $\{u, v, uv\} \subseteq H_{k+1}$. Thus \mathbf{H}_{k+1} is a partial *ComIdBass*-groupoid. \square

Proposition 2.3. (a) $H_k \subset H_{k+1}$, (b) $D_{k+1} \subset H_{k+1}^2$. \square

Proposition 2.4. If $\mathbf{G} \in \text{ComIdBass}$ and $\lambda : B \rightarrow G$, then for each $k \geq 0$, there is a partial homomorphism $\varphi_{k+1} : H_{k+1} \rightarrow G$, such that φ_k is the restriction of φ_{k+1} on H_k and $\varphi_0 = \lambda$. \square

Theorem 2. Let $H = \cup\{H_k \mid k \geq 0\}$. Then $\mathbf{H} = (H, \cdot)$ is a free *ComIdBass*-groupoid with the basis B .

Proof. In the same way as in Theorem 1, one can prove that $\mathbf{H} \in \text{ComIdBass}$, it is generated by B and if $\mathbf{G} \in \text{ComIdBass}$ and $\lambda : B \rightarrow G$ is a mapping, then $\varphi = \cup_{k \geq 0} \varphi_k : H \rightarrow G$ is the homomorphic extension of λ . \square

Remark 6. For the construction of a free object in the variety *IdBass* with a basis B , a theorem similar to Theorem 2 can be used. Then the construction is essentially the same as for *ComIdBass*, except for that here the free idempotent semigroup $\{a, b, ab, ba, aba, bab\}$ generated by $\{a, b\}$ is used, instead of a free commutative idempotent semigroup $\{a, b, ab\}$ generated by $\{a, b\}$.

References

- [1] Bruck, R. H., A Survey on Binary Systems. Berlin, Göttingen, Heidelberg: Springer-Verlag 1958.
- [2] Maljcev, A. I., Algebraicheskie sistemi. Moskva: Nauka 1970 (in Russian).
- [3] Čupona, Ć., Celakoski, N., Ilić, S., On monoassociative groupoids. Matematički bilten 26 (LII) (2002), 5-16.

Received by the editors July 9, 2003